

## Referenceliste for NKR for forebyggelse af fald hos ældre.

Høringsudgave, samlet litteratur fra Indledning/baggrund samt inkluderet referencer. (060717/BHP)

- (1) Abrahamowicz M, Bartlett G, Tamblyn R, du Berger R. Modeling cumulative dose and exposure duration provided insights regarding the associations between benzodiazepines and injuries. *J Clin Epidemiol* 2006;59(4):393-403.
- (2) Ahearn DJ, Umapathy D. Vestibular impairment in older people frequently contributes to dizziness as part of a geriatric syndrome. *Clin Med (Lond)* 2015;15(1):25-30.
- (3) Airagnes G, Pelissolo A, Lavallee M, Flament M, Limosin F. Benzodiazepine Misuse in the Elderly: Risk Factors, Consequences, and Management. *Curr Psychiatry Rep* 2016;18(10):89-016-0727-9.
- (4) Arantes P.M.M., Dias J.M.D., Fonseca F.F., et al. . Effect of a Program Based on Balance Exercises on Gait, Functional Mobility, Fear of Falling, and Falls in Pre frail Older Women: A Randomized Clinical Trial. Opdateret: 2015. Senest hentet: 2.31.
- (5) Bakken MS, Engeland A, Engesaeter LB, Ranhoff AH, Hunskaar S, Ruths S. Risk of hip fracture among older people using anxiolytic and hypnotic drugs: a nationwide prospective cohort study. *Eur J Clin Pharmacol* 2014;70(7):873-80.
- (6) Barker AL, Morello RT, Wolfe R, Brand CA, Haines TP, Hill KD, et al. 6-PACK programme to decrease fall injuries in acute hospitals: cluster randomised controlled trial. *BMJ* 2016;352:h6781.
- (7) Bloch F, Thibaud M, Dugue B, Breque C, Rigaud AS, Kemoun G. Psychotropic drugs and falls in the elderly people: updated literature review and meta-analysis (Structured abstract). 2011;23(2):329-346.
- (8) Chu M.M.-L., Fong K.N.-K., Lit A.C.-H., et al. . An Occupational Therapy Fall Reduction Home Visit Program for Community-Dwelling Older Adults in Hong Kong After an Emergency Department Visit for a Fall. Opdateret: 2016.
- (9) Clemson L, Mackenzie L, Ballinger C, Close JCT, Cumming RG. Environmental interventions to prevent falls in community-dwelling older people: a meta-analysis of randomized trials. *J Aging Health* 2008;20(8):954-71.
- (10) COOKSEY FS. Rehabilitation in vestibular injuries. *Proc R Soc Lond B Biol Sci* 1946;39:273-278.
- (11) Cox H, Puffer S, Morton V, Cooper C, Hodson J, Masud T, et al. Educating nursing home staff on fracture prevention: a cluster randomised trial. *Age Ageing* 2008;37(2):167-72.
- (12) Cumming RG, Thomas M, Szonyi G, Salkeld G, O'Neill E, Westbury C, et al. Home visits by an occupational therapist for assessment and modification of environmental hazards: a randomized trial of falls prevention. *J Am Geriatr Soc* 1999;47(12):1397-1402.
- (13) Danmarks Statistik. 65+ et portræt af de ældres liv, arbejdsliv og sociale situation Danmarks Statistik, 2012. 119 s. (TemaPubl 2012:3).
- (14) Day L, Hill KD, Stathakis VZ, Flicker L, Segal L, Cicuttini F, et al. Impact of tai-chi on falls among preclinically disabled older people. A randomized controlled trial. *Journal of the American Medical Directors Association* 2015;16(5):420-6.
- (15) Enticott JC, Vitkovic JJ, Reid B, O'Neill P, Paine M. Vestibular rehabilitation in individuals with inner-ear dysfunction: a pilot study. *Audiol Neurotol* 2008;13(1):19-28.

- (16) Feldman F, Chaudhury H. Falls and the physical environment: a review and a new multifactorial falls-risk conceptual framework. *Can J Occup Ther* 2008;75(2):82-95.
- (17) Fernandez-Arguelles E.L., Rodriguez-Mansilla J., Antunez L.E., Garrido-Ardila E.M., Munoz R.P. Effects of dancing on the risk of falling related factors of healthy older adults: A systematic review. *Arch Gerontol Geriatr* 2015;60(1):1-8.
- (18) Fitzharris MP, Day L, Lord SR, Gordon I, Fildes B. The Whitehorse NoFalls trial: effects on fall rates and injurious fall rates. *Age Ageing* 2010;39(6):728-33.
- (19) Flint A.J., Iaboni A., Mulsant B.H., Rothschild A.J., Whyte E.M. and Meyers B.S. . Sertraline on risk of falling in older adults with psychotic depression on olanzapine: Results of a randomized placebo-controlled trial. *Opdateret*: 2014. Senest hentet: 4.22.
- (20) Gebara MA, Lipsey KL, Karp JF, Nash MC, Iaboni A, Lenze EJ. Cause or Effect? Selective Serotonin Reuptake Inhibitors and Falls in Older Adults: A Systematic Review. *Am J Geriatr Psychiatry* 2015;23(10):1016-28.
- (21) Gillespie L.D., Robertson M.C., Gillespie W.J., Sherrington C., Gates S., Clemson L.M., et al. Interventions for preventing falls in older people living in the community. *Cochrane Database Syst Rev* 2012;9:CD007146.
- (22) Giray M, Kirazli Y, Karapolat H, Celebisoy N, Bilgen C, Kirazli T. Short-term effects of vestibular rehabilitation in patients with chronic unilateral vestibular dysfunction: a randomized controlled study. *Arch Phys Med Rehabil* 2009;90(8):1325-31.
- (23) Golden AG, Ma Q, Nair V, Florez HJ, Roos BA. Risk for fractures with centrally acting muscle relaxants: an analysis of a national Medicare Advantage claims database. *Ann Pharmacother* 2010;44(9):1369-75.
- (24) Granacher U, Muehlbauer T, Bridenbaugh SA, Wolf M, Roth R, Gschwind Y, et al. Effects of a salsa dance training on balance and strength performance in older adults. *Gerontology* 2012;58(4):305-12.
- (25) Halvarsson A, Franzen E, Faren E, Olsson E, Oddsson L, Stahle A. Long-term effects of new progressive group balance training for elderly people with increased risk of falling - a randomized controlled trial. *Clin Rehabil* 2013;27(5):450-8.
- (26) Halvarsson A, Franzen E, Stahle A. Balance training with multi-task exercises improves fall-related self-efficacy, gait, balance performance and physical function in older adults with osteoporosis: a randomized controlled trial. *Clin Rehabil* 2015;29(4):365-75.
- (27) Ham AC, Ziere G, Broer L, Swart KMA, Enneman AW, van Dijk S.C., et al. CYP2C9 Genotypes Modify Benzodiazepine-Related Fall Risk: Original Results From Three Studies With Meta-Analysis. *J AM MED DIR ASSOC* 2017;18(1):88.e1-88.e15.
- (28) Harun A, Li C, Bridges JF, Agrawal Y. Understanding the Experience of Age-Related Vestibular Loss in Older Individuals: A Qualitative Study. *Patient* 2016;9(4):303-309.
- (29) Herdman S.J. (editor). Assessment and interventions for the patient with complete vestibular loss. 3rd edn. Assessment and interventions for the patient with complete vestibular loss : F.A. Davis, Philadelphia, 2007.
- (30) Hill KD, Wee R. Psychotropic Drug-Induced Falls in Older People. *Drugs Aging* 2012;29(1):15-30.
- (31) Hirase T, Inokuchi S, Matsusaka N, Okita M. Effects of a balance training program using a foam rubber pad in community-based older adults: a randomized controlled trial. *Journal of geriatric physical therapy* (2001) 2015;38(2):62-70.

- (32) Hoffer ME, Balaban CD. Vestibular rehabilitation: ready for the mainstream. *NeuroRehabilitation* 2011;29(2):125-2011-0686.
- (33) Hødrup S, Sorensen TI, Gronbaek M, Schroll M. Incidence and characteristics of falls leading to hospital treatment: a one-year population surveillance study of the Danish population aged 45 years and over. *Scand J Public Health* 2003;31(1):24-30.
- (34) Horak FB, Jones-Rycewicz C, Black FO, Shumway-Cook A. Effects of vestibular rehabilitation on dizziness and imbalance. *Otolaryngol Head Neck Surg* 1992;106(2):175-180.
- (35) Howe TE, Rochester L, Jackson A, Banks PM, Blair VA. Exercise for improving balance in older people. *Cochrane Database Syst Rev* 2007;(4)(4):CD004963.
- (36) Institute of Medicine (US) Division of Health Promotion and Disease Prevention. The Second Fifty Years: Promoting Health and Preventing Disability. Washington (DC) : National Academy of Sciences, 1992.
- (37) Jakobsen JC, Katakam KK, Schou A, Hellmuth SG, Stallknecht SE, Leth-Møller K, et al. Selective serotonin reuptake inhibitors versus placebo in patients with major depressive disorder. A systematic review with meta-analysis and Trial Sequential Analysis. *BMC Psychiatry* 2017;17(1):58-016-1173-2.
- (38) Kamei T, Kajii F, Yamamoto Y, Irie Y, Kozakai R, Sugimoto T, et al. Effectiveness of a home hazard modification program for reducing falls in urban community-dwelling older adults: A randomized controlled trial. *Japan journal of nursing science : JJNS* 2015;12(3):184-97.
- (39) Kannus P, Sievanen H, Palvanen M, Jarvinen T, Parkkari J. Prevention of falls and consequent injuries in elderly people. *Lancet* 2005;366(9500):1885-1893.
- (40) Karinkanta S, Nupponen R, Heinonen A, Pasanen M, Sievanen H, Uusi-Rasi K, et al. Effects of exercise on health-related quality of life and fear of falling in home-dwelling older women. *J Aging Phys Act* 2012;20(2):198-214.
- (41) Keall MD, Piers N, Howden-Chapman P, Cunningham C, Cunningham M, Guria J, et al. Home modifications to reduce injuries from falls in the home injury prevention intervention (HIPI) study: a cluster-randomised controlled trial. *Lancet (London, England)* 2015;385(9964):231-8.
- (42) Kelsey JL, Berry SD, Procter-Gray E, Quach L, Nguyen US, Li W, et al. Indoor and outdoor falls in older adults are different: the maintenance of balance, independent living, intellect, and Zest in the Elderly of Boston Study. *J Am Geriatr Soc* 2010;58(11):2135-2141.
- (43) Korpelainen R, Keinanen-Kiukkaanniemi S, Heikkinen J, Vaananen K, Korpelainen J. Effect of impact exercise on bone mineral density in elderly women with low BMD: a population-based randomized controlled 30-month intervention. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA* 2006;17(1):109-18.
- (44) Krampe J. . Exploring the Effects of Dance-Based Therapy on Balance and Mobility in Older Adults. *Opdateret: 2013. Senest hentet: 1.35.*
- (45) Lawton B., Rose S.B., Elley C.R., Garrett S., Dowell A., Fenton A., et al. Prescribing exercise to improve quality of life? *Climacteric* 2011;14:77.
- (46) Leung DPK, Chan CKL, Tsang HWH, Tsang WWN, Jones AYM. Tai chi as an intervention to improve balance and reduce falls in older adults: A systematic and meta-analytical review. *Altern Ther Health Med* 2011;17(1):40-8.

- (47) Lin M, Hwang H, Wang Y, Chang S, Wolf SL. Community-based tai chi and its effect on injurious falls, balance, gait, and fear of falling in older people. *Phys Ther* 2006;86(9):1189-201.
- (48) Liston MB, Bamiou DE, Martin F, Hopper A, Koohi N, Luxon L, et al. Peripheral vestibular dysfunction is prevalent in older adults experiencing multiple non-syncopal falls versus age-matched non-fallers: a pilot study. *Age Ageing* 2014;43(1):38-43.
- (49) Logghe IHJ, Zeeuwe PEM, Verhagen AP, Wijnen-Sponselee R, Willemsen SP, Bierma-Zeinstra S, et al. Lack of effect of Tai Chi Chuan in preventing falls in elderly people living at home: a randomized clinical trial. *J Am Geriatr Soc* 2009;57(1):70-5.
- (50) Madureira MM, Bonfa E, Takayama L, Pereira RMR. A 12-month randomized controlled trial of balance training in elderly women with osteoporosis: improvement of quality of life. *Maturitas* 2010;66(2):206-11.
- (51) McDonnell MN, Hillier SL. Vestibular rehabilitation for unilateral peripheral vestibular dysfunction. *Cochrane Database Syst Rev* 2015;1:CD005397.
- (52) Melzer I, Oddsson LI. Improving balance control and self-reported lower extremity function in community-dwelling older adults: a randomized control trial. *Clin Rehabil* 2013;27(3):195-206.
- (53) Merom D., Mathieu E., Cerin E., et al. . Social Dancing and Incidence of Falls in Older Adults: A Cluster Randomised Controlled Trial. *Opdateret: 2016. Senest hentet: 8.13.*
- (54) Møller H, Damm M, Laursen B. Ulykkeri Danmarks 1990-2009 Statens Institut for Folkesundhed, Syddansk Universitet, 2012.
- (55) Muir SW, Berg K, Chesworth B, Klar N, Speechley M. Quantifying the magnitude of risk for balance impairment on falls in community-dwelling older adults: a systematic review and meta-analysis. *J Clin Epidemiol* 2010;63(4):389-406.
- (56) Nelson J.C., Oakes T.M.M., Liu P., et al. . Assessment of falls in older patients treated with duloxetine: A secondary analysis of a 24-week randomized, placebo-controlled trial. *Opdateret: 2013. Senest hentet: 1.15.*
- (57) Nevitt MC, Cummings SR, Kidd S, Black D. Risk factors for recurrent nonsyncopal falls. A prospective study. *JAMA* 1989;261(18):2663-2668.
- (58) Nguyen MH, Kruse A. A randomized controlled trial of Tai chi for balance, sleep quality and cognitive performance in elderly Vietnamese. *Clinical interventions in aging* 2012;7:185-90.
- (59) Nick N, Petramfar P, Ghodsbini F, Keshavarzi S, Jahanbin I. The Effect of Yoga on Balance and Fear of Falling in Older Adults. *PM & R : the journal of injury, function, and rehabilitation* 2016;8(2):145-51.
- (60) Nurminen J, Puustinen J, Lahteenmaki R, Vahlberg T, Lyles A, Partinen M, et al. Handgrip strength and balance in older adults following withdrawal from long-term use of temazepam, zopiclone or zolpidem as hypnotics. *BMC Geriatr* 2014;14:121-2318-14-121.
- (61) Nyberg L, Gustafson Y, Janson A, Sandman PO, Eriksson S. Incidence of falls in three different types of geriatric care. A Swedish prospective study. *Scand J Soc Med* 1997;25(1):8-13.
- (62) Oderda LH, Young JR, Asche CV, Pepper GA. Psychotropic-related hip fractures: meta-analysis of first-generation and second-generation antidepressant and antipsychotic drugs. *Ann Pharmacother* 2012;46(7-8):917-28.

- (63) O'Loughlin JL, Robitaille Y, Boivin JF, Suissa S. Incidence of and risk factors for falls and injurious falls among the community-dwelling elderly. *Am J Epidemiol* 1993;137(3):342-354.
- (64) Orr R. Contribution of muscle weakness to postural instability in the elderly. A systematic review. *Eur J Phys Rehabil Med* 2010;46(2):183-220.
- (65) Park H, Satoh H, Miki A, Urushihara H, Sawada Y. Medications associated with falls in older people: systematic review of publications from a recent 5-year period. *Eur J Clin Pharmacol* 2015;71(12):1429-40.
- (66) Peel NM. Epidemiology of falls in older age. *Can J Aging* 2011;30(1):7-19.
- (67) Peterson MD, Rhea MR, Sen A, Gordon PM. Resistance exercise for muscular strength in older adults: a meta-analysis. *Ageing Res Rev* 2010;9(3):226-237.
- (68) Pighills AC, Torgerson DJ, Sheldon TA, Drummond AE, Bland JM. Environmental assessment and modification to prevent falls in older people. *J Am Geriatr Soc* 2011;59(1):26-33.
- (69) Pliske G., Emmermacher P., Weinbeer V. and Witte K. . Changes in dual-task performance after 5 months of karate and fitness training for older adults to enhance fall prevention. *Opdateret: 2015*.
- (70) Prudham D, Evans JG. Factors associated with falls in the elderly: a community study. *Age Ageing* 1981;10(3):141-146.
- (71) R. Rosenberg, K. S. Christensen and M. B. Jørgensen. . Benzodiazepiner (anxiolytika) N05BA. *Opdateret: Revideret: 27.02.2017. Senest hentet: 3. juli 2017.2017.*  
Link: <https://pro.medicin.dk/Laegemiddelgrupper/Grupper/239010>.
- (72) Rossat A, Fantino B, Bongue B, Colvez A, Nitenberg C, Annweiler C, et al. Association between benzodiazepines and recurrent falls: a cross-sectional elderly population-based study. *J Nutr Health Aging* 2011;15(1):72-7.
- (73) Sakamoto K, Endo N, Harada A, Sakada T, Tsushita K, Kita K, et al. Why not use your own body weight to prevent falls? A randomized, controlled trial of balance therapy to prevent falls and fractures for elderly people who can stand on one leg for <15 s. *J Orthop Sci* 2013;18(1):110-20.
- (74) Salonoja M, Salminen M, Vahlberg T, Aarnio P, Kivela SL. Withdrawal of psychotropic drugs decreases the risk of falls requiring treatment. *Arch Gerontol Geriatr* 2012;54(1):160-167.
- (75) Seo B.D., Kim B.J. and Singh K. . The comparison of resistance and balance exercise on balance and falls efficacy in older females. *Opdateret: 2012. Senest hentet: 5.3.*
- (76) Serra-Rexach J, Bustamante-Ara N, Hierro Villaran M, Gonzalez Gil P, Sanz Ibanez M,J., Blanco Sanz N, et al. Short-term, light- to moderate-intensity exercise training improves leg muscle strength in the oldest old: a randomized controlled trial. *J Am Geriatr Soc* 2011;59(4):594-602.
- (77) Sheffield C, Smith CA, Becker M. Evaluation of an agency-based occupational therapy intervention to facilitate aging in place. *Gerontologist* 2013;53(6):907-18.
- (78) Sherrington C, Tiedemann A, Fairhall N, Close JC, Lord SR. Exercise to prevent falls in older adults: an updated meta-analysis and best practice recommendations. *N S W Public Health Bull* 2011;22(3-4):78-83.
- (79) Shigematsu R, Okura T, Nakagaichi M, Tanaka K, Sakai T, Kitazumi S, et al. Square-stepping exercise and fall risk factors in older adults: a single-blind, randomized controlled trial. *The journals of gerontology Series A, Biological sciences and medical sciences* 2008;63(1):76-82.

- (80) Silva RB, Eslick GD, Duque G. Exercise for falls and fracture prevention in long term care facilities: a systematic review and meta-analysis. *J AM MED DIR ASSOC* 2013;14(9):685-9.e2.
- (81) Smith-Wheelock M, Shepard NT, Telian SA. Physical therapy program for vestibular rehabilitation. *Am J Otol* 1991;12(3):218-225.
- (82) Sousa N, Mendes R. Comparison of effects of resistance and multicomponent training on falls prevention in institutionalized elderly women. *J Am Geriatr Soc* 2015;63(2):396-7.
- (83) Sundhedsstyrelsen. National klinisk retningslinje for Ernærings- og træningsindsatser til ældre med geriatriske problemstillinger. [version 1.0]. Sundhedsstyrelsen, december 2016.
- (84) Sundhedsstyrelsen. Vejledning om ordination af afhængighedsskabende lægemidler og om substitutionsbehandling af personer med opioidafhængighed. Version 1.0. Sundhedsstyrelsen, 8. juni 2007.
- (85) Sundhedsstyrelsen. Faldpatienter i den kliniske hverdag - rådgivning fra Sundhedsstyrelsen Sundhedsstyrelsen, 2006.
- (86) Tiedemann A, O'Rourke S, Sesto R, Sherrington C. A 12-week Iyengar yoga program improved balance and mobility in older community-dwelling people: a pilot randomized controlled trial. *The journals of gerontology Series A, Biological sciences and medical sciences* 2013;68(9):1068-75.
- (87) Tinetti ME, Speechley M, Ginter SF. Risk factors for falls among elderly persons living in the community. *N Engl J Med* 1988;319(26):1701-1707.
- (88) Trombetti A., Hars M., Herrmann F.R., Kressig R.W., Ferrari S. and Rizzoli R. . Effect of music-based multitask training on gait, balance, and fall risk in elderly people: A randomized controlled trial. Opdateret: 2011. Senest hentet: 6.171.
- (89) Turner S, Arthur G, Lyons RA, Weightman AL, Mann MK, Jones SJ, et al. Modification of the home environment for the reduction of injuries. *Cochrane Database Syst Rev* 2011(2):CD003600.
- (90) Ullmann G, Williams HG, Hussey J, Durstine JL, McClenaghan BA. Effects of Feldenkrais exercises on balance, mobility, balance confidence, and gait performance in community-dwelling adults age 65 and older. *Journal of alternative and complementary medicine (New York, N Y )* 2010;16(1):97-105.
- (91) Uzun S, Kozumplik O, Jakovljevic M, Sedici B. Side effects of treatment with benzodiazepines. *Psychiatr Danub* 2010;22(1):90-93.
- (92) Vaapio S., Puustinen J., Salminen M.J., et al. . Symptoms associated with long-term benzodiazepine use in elderly individuals aged 65 years and older: A longitudinal descriptive study. Opdateret: 2015. Senest hentet: 1.9.
- (93) Van der Hooft C. Schoofs M.W.C.J., Ziere G., et al. . Inappropriate benzodiazepine use in older adults and the risk of fracture. Opdateret: 2008. Senest hentet: 2.66.
- (94) van Gaal B,G.I., Schoonhoven L, Mintjes JAJ, Borm GF, Hulscher MEJL, Defloor T, et al. Fewer adverse events as a result of the SAFE or SORRY? programme in hospitals and nursing homes. part i: primary outcome of a cluster randomised trial. *Int J Nurs Stud* 2011;48(9):1040-8.
- (95) van Gaal B,G.I., Schoonhoven L, Vloet LCM, Mintjes JAJ, Borm GF, Koopmans RTCM, et al. The effect of the SAFE or SORRY? programme on patient safety knowledge of nurses in hospitals and nursing homes: a cluster randomised trial. *Int J Nurs Stud* 2010;47(9):1117-25.

- (96) Vereeck L, Wuyts FL, Truijen S, De Valck C, Van dH. The effect of early customized vestibular rehabilitation on balance after acoustic neuroma resection. *Clin Rehabil* 2008;22(8):698-713.
- (97) Vikman I, Nordlund A, Naslund A, Nyberg L. Incidence and seasonality of falls amongst old people receiving home help services in a municipality in northern Sweden. *Int J Circumpolar Health* 2011;70(2):195-204.
- (98) Voukelatos A, Cumming RG, Lord SR, Rissel C. A randomized, controlled trial of tai chi for the prevention of falls: the Central Sydney tai chi trial. *J Am Geriatr Soc* 2007;55(8):1185-91.
- (99) Voyer P, Preville M, Cohen D, Berbiche D, Beland SG. The prevalence of benzodiazepine dependence among community-dwelling older adult users in Quebec according to typical and atypical criteria. *Can J Aging* 2010;29(2):205-213.
- (100) Warden SJ, Fuchs RK. Do Selective Serotonin Reuptake Inhibitors (SSRIs) Cause Fractures? *Curr osteoporos rep* 2016;14(5):211-8.
- (101) Wilkinson P, Izmeth Z. Continuation and maintenance treatments for depression in older people. 2016(9).
- (102) Woo J, Hong A, Lau E, Lynn H. A randomised controlled trial of Tai Chi and resistance exercise on bone health, muscle strength and balance in community-living elderly people. *Age Ageing* 2007;36(3):262-8.
- (103) Yardley L. and Kirby S. . Evaluation of booklet-based self-management of symptoms in Meniere disease: A randomized controlled trial. *Opdateret: 2006. Senest hentet: 5.68.*
- (104) Yardley L, Donovan-Hall M, Francis K, Todd C. Older people's views of advice about falls prevention: a qualitative study. *Health Educ Res* 2006;21(4):508-517.
- (105) Youkhana S., Dean C.M., Wolff M., Sherrington C. and Tiedemann A. . Yoga-based exercise improves balance and mobility in people aged 60 and over: A systematic review and meta-analysis. *Opdateret: 2016. Senest hentet: 1.45.*
- (106) Zhang J, Ishikawa-Takata K, Yamazaki H, Morita T, Ohta T. The effects of Tai Chi Chuan on physiological function and fear of falling in the less robust elderly: an intervention study for preventing falls. *Arch Gerontol Geriatr* 2006;42(2):107-16.