

T-2 and HT-2 TOXINS

The provisions in this document have not been adopted or endorsed by the European Commission. Any views expressed are the preliminary views of the Commission services and may not in any circumstances be regarded as stating an official position of the Commission.

SUGGESTED POSSIBLE MAXIMUM LEVELS FOR T-2 and HT-2 TOXINS AS DISCUSSED IN THE WORKING GROUP “AGRICULTURAL CONTAMINANTS” IN VIEW OF THIS TARGETED STAKEHOLDER CONSULTATION

Food	Suggested levels for the sum of T-2 and HT-2 toxin in µg/kg – for discussion
1. Unprocessed cereals	
1.1. barley, maize and durum wheat ⁽¹⁸⁾	100
1.2. oats (with husk)	1000
1.3. wheat, rye and other cereals ⁽¹⁸⁾	50
2. Cereal grains placed on the market for the final consumer	
2.1. oats, maize, durum wheat	50
2.2. other cereals	20
3. Cereal products	
3.1. cereal bran, oat milling products and maize milling products	50
3.2. other cereal milling products	20
3.3. bread (including small bakery wares), pastries, biscuits, cereal snacks, breakfast cereals including formed cereal flakes, pasta	Option A: 20 Option B: ML applicable to these products is the ML determined in application of Article 2 of Regulation (EC) 1881/2006
3.4. cereal based foods for infants and young children	10