



1 PICO 1 - Medicin + non-farmakologiske interventioner vs waitlist/TAU - Primæranalyser


1.1 Funktionsniveau - observatørvurderet - behandlingsafslutning - SMD




1.2 Funktionsniveau - observatørvurderet - behandlingsafslutning - CGI severity




1.3 Funktionsniveau - observatørvurderet - længste opfølgning - SMD



1.4 Funktionsniveau - observatørvurderet - længste opfølgning - CGI severity



1.5 Funktionsniveau - selvvrderet - behandlingsafslutning - SMD



Footnotes

- (1) Work and Social Adjustment Scale (WSAS). Skala 0-8, lower is better.
(2) BRIEF-A = Behavior Rating Inventory of Executive Function-Adult Version, skala 1-3. 75 item, lower is better
(3) Canadian Occupational Performance Measure (COPM), range: 1-10, higher is better (derfor -)
(4) BRIEF-A Behavior Rating Inventory of Executive Function - Adult Version, total sum score, lower is better

- (5) BRIEF-A = Behavior Rating Inventory of Executive Functioning—Adult Version, range 1-3, lower is better
(6) BRIEF-ASR = Behavior Rating Inventory of Executive Function-Adult Self-Report Version, lower is better
(7) BDEFS-SF: Barkley Deficits in Executive Functioning Scale - short form, range 20-80, lower is better
(8) BDEFS-SF = Barkley Deficits in Executive Functioning Scale-short form, lower is better


- (E) Incomplete outcome data (attrition bias)
(F) Selective reporting (reporting bias)
(G) Other bias

1.6 Funktionsniveau - selvurderet - længste opfølgning


Study or Subgroup	Non-farm. interventioner		Waitlist/TAU		Weight	Mean Difference		Mean Difference	Risk of Bias						
	Mean	SD	Total	Mean	SD	IV, Random, 95% CI	IV, Random, 95% CI	IV, Random, 95% CI	A	B	C	D	E	F	G
1.6.1 Kognitiv adfærdsterapi															
Dittner 2018 (1)	10.2	6.9	27	16.8	8.8	18 100.0%	-6.60 [-11.43, -1.77]								
Subtotal (95% CI)			27			18 100.0%	-6.60 [-11.43, -1.77]								
Heterogeneity: Not applicable															
Test for overall effect: Z = 2.68 (P = 0.007)															
Total (95% CI)			27			18 100.0%	-6.60 [-11.43, -1.77]								
Heterogeneity: Not applicable															
Test for overall effect: Z = 2.68 (P = 0.007)															
Test for subgroup differences: Not applicable															

Footnotes


- (1) Work and Social Adjustment Scale (WSAS). Skala 0-8 på 5 items. Lower is better. 43 ugers opfølgning.
Risk of bias legend
(A) Random sequence generation (selection bias)
(B) Allocation concealment (selection bias)
(C) Blinding of participants and personnel (performance bias)
(D) Blinding of outcome assessment (detection bias)
(E) Incomplete outcome data (attrition bias)
(F) Selective reporting (reporting bias)
(G) Other bias




1.7 Alvorlige hændelser - længste opfølgning



1.8 Livskvalitet - behandlingsafslutning - SMD




1.9 ADHD symptomer - observatørvurderet - behandlingsafslutning - SMD



Test for overall effect: $Z = 3.94$ ($P < 0.0001$)
Test for subgroup differences: $\text{Chi}^2 = 35.35$, $\text{df} = 5$ ($P < 0.00001$), $I^2 = 85.9\%$


Footnotes

(1) Adult Barkley Current Symptoms Scale (CSS). Skala 0-3. totalscore max 54 point. Lower is better
(2) The Barkley ADHD Current Symptoms Scale (BCS), 0-54. lower is better
(3) Adult ADHD Self-Report Scale version 1.1 (ASRS-v1.1), range: 0-72, lower is better
(4) ADHD rating scale, Range 0-54 Lower is better
(5) ASRS Adult ADHD Self-Report Scale, total sum score, 0-72, lower is better
(6) CAARS (long version) total score, range: 0-198, lower is better
(7) CAARS index = Conners' Adult ADHD Self-Rating Scale, ADHD index, lower is better
(8) Adult ADHD Self-Report Scale version 1.1 (ASRS-v1.1), range: 0-72, lower is better
(9) CAARS investigator rating version, lower is better
(10) 10-point category scale based on the DSM IV diagnostic criteria for ADHD, range? Lower is better




1.10 ADHD symptomer - selvvurderet - behandlingsafslutning - SMD

Study or Subgroup	Non-farm. interventioner		Waitlist/TAU		Weight	Std. Mean Difference		Std. Mean Difference		Risk of Bias						
	Mean	SD	Total	Mean	SD	Total	Mean	IV, Random, 95% CI	IV, Random, 95% CI	A	B	C	D	E	F	G
1.10.1 Kognitiv adfærdsterapi																
Huang 2019 (1)	17.85	8	33	24.44	4.76	16	8.0%	-0.91 [-1.54, -0.28]								
Pan 2022 (2)	20.05	9.35	49	20.75	9.3	49	12.8%	-0.07 [-0.47, 0.32]								
Petterson (CBT) 2017 (3)	27.31	12.28	13	29.72	8.17	18	6.7%	-0.23 [-0.95, 0.48]								
Safren 2005 (4)	14.75	8.65	16	23.87	9.92	15	6.3%	-0.96 [-1.71, -0.21]								
Virta (CBT) 2010 (5)	45.4	11.7	10	47.8	12.8	5	3.6%	-0.19 [-1.26, 0.89]								
Subtotal (95% CI)			121			103	37.5%	-0.45 [-0.86, -0.04]								
Heterogeneity: $\tau^2 = 0.10$; $\text{Chi}^2 = 7.60$, $\text{df} = 4$ ($P = 0.11$); $I^2 = 47\%$																
Test for overall effect: $Z = 2.18$ ($P = 0.03$)																
1.10.2 Dialektisk adfærdsterapi																
Moritz 2021 (6)	30.77	8.228	13	30.79	7.934	14	6.2%	-0.00 [-0.76, 0.75]								
Subtotal (95% CI)			13			14	6.2%	-0.00 [-0.76, 0.75]								
Heterogeneity: Not applicable																
Test for overall effect: $Z = 0.01$ ($P = 1.00$)																
1.10.3 Dialektisk adfærdsterapi/kognitiv adfærdsterapi																
Philipsen (methylphenidat) 2015 (7)	81.3	32.4	87	81.7	34.1	94	15.7%	-0.01 [-0.30, 0.28]								
Subtotal (95% CI)			87			94	15.7%	-0.01 [-0.30, 0.28]								
Heterogeneity: Not applicable																
Test for overall effect: $Z = 0.08$ ($P = 0.94$)																
1.10.5 Mindfulness-baseret kognitiv terapi																
Hepark 2019 (8)	23	7.3	41	28.8	6.9	42	11.5%	-0.81 [-1.26, -0.36]								
Subtotal (95% CI)			41			42	11.5%	-0.81 [-1.26, -0.36]								
Heterogeneity: Not applicable																
Test for overall effect: $Z = 3.54$ ($P = 0.0004$)																
1.10.6 Kognitiv træning/psykosociale interventioner																
Virta (CT) 2010 (9)	41.2	4.9	9	47.8	12.8	5	3.3%	-0.74 [-1.87, 0.40]								
Subtotal (95% CI)			9			5	3.3%	-0.74 [-1.87, 0.40]								
Heterogeneity: Not applicable																
Test for overall effect: $Z = 1.26$ ($P = 0.21$)																
1.10.7 Træning af arbejdshukommelse																
Liu 2017 (10)	46.85	8.81	34	47.21	9.84	34	10.9%	-0.04 [-0.51, 0.44]								
Mawjee 2015 (11)	46.48	9.89	32	47.25	11.51	32	10.6%	-0.07 [-0.56, 0.42]								
Mawjee 2017 (12)	46.57	8.44	8	51.44	11.23	9	4.3%	-0.46 [-1.43, 0.51]								
Subtotal (95% CI)			74			75	25.8%	-0.10 [-0.42, 0.22]								
Heterogeneity: $\tau^2 = 0.00$; $\text{Chi}^2 = 0.61$, $\text{df} = 2$ ($P = 0.74$); $I^2 = 0\%$																
Test for overall effect: $Z = 0.60$ ($P = 0.55$)																



1.11 ADHD symptom - uopmærksomhed - observatørvurderet - behandlingsafslutning - SMD




Footnotes


(1) CAARS-Inattention/memory subscale, range?, lower is better

(2) BARKLEY ADULT ADHD RATING SCALE-IV (BAARS-IV); Inattention, range 0-27, lower is better


1.12 ADHD symptom - uopmærksomhed - selvvrurderet - behandlingsafslutning - SMD




1.13 ADHD symptom - hyperaktiv impulsivitet - observatørvurderet - behandlingsafslutning



1.15 Frafald - behandlingsafslutning






1.16 Misbrug og afhængighed - længste opfølgning (0 RCT)

Study or Subgroup	Non-farm. interventioner		Waitlist/TAU		Total Weight	Mean Difference		Risk of Bias						
	Mean	SD	Total	Mean	SD	IV, Random	95% CI	A	B	C	D	E	F	G
Total (95% CI)			0			0								
Heterogeneity:	Not applicable													
Test for overall effect:	Not applicable													

Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias



1.17 Fravær fra arbejde - behandlingsafslutning (0 RCT)

Study or Subgroup	Non-farm. interventioner		Waitlist/TAU		Total Weight	Risk Ratio		Risk Ratio	Risk of Bias						
	Events	Total	Events	Total		M-H, Random, 95% CI	M-H, Random, 95% CI		A	B	C	D	E	F	G

Total (95% CI)

0000

Not estimable

Total events

0

0

Heterogeneity: Not applicable

Test for overall effect: Not applicable



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

1.18 Kriminalitet - længste opfølgning (0 RCT)

Study or Subgroup	Non-farm. interventioner		Waitlist/TAU		Total Weight	Risk Ratio		Risk Ratio		Risk of Bias						
	Events	Total	Events	Total		M-H, Random, 95% CI	M-H, Random, 95% CI	M-H, Random, 95% CI	M-H, Random, 95% CI	A	B	C	D	E	F	G

Total (95% CI)

Total events

Heterogeneity: Not applicable

Test for overall effect: Not applicable

Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias



1.19 Ophor af medicinsk behandling - behandlingsafslutning

Study or Subgroup	Non-farm. interventioner		Waitlist/TAU		Total Weight	Risk Ratio		Risk Ratio M-H, Random, 95% CI	Risk of Bias						
	Events	Total	Events	Total		M-H, Random, 95% CI	95% CI		A	B	C	D	E	F	G
Pan 2022	0	49	1	49	100.0%	0.33	[0.01, 7.99]								
Total (95% CI)		49		49	100.0%	0.33	[0.01, 7.99]								
Total events	0		1												
Heterogeneity: Not applicable															
Test for overall effect: Z = 0.68 (P = 0.50)															



1.20 Ophør af medicinsk behandling - længste follow-up

Study or Subgroup	Non-farm. interventioner		Waitlist/TAU		Total Weight	Risk Ratio		Risk Ratio	Risk of Bias						
	Events	Total	Events	Total		M-H, Random, 95% CI	M-H, Random, 95% CI		A	B	C	D	E	F	G
Pan 2022	5	49	2	49	100.0%		2.50 [0.51, 12.28]								
Total (95% CI)		49		49	100.0%		2.50 [0.51, 12.28]								
Total events	5		2												
Heterogeneity: Not applicable															
Test for overall effect: Z = 1.13 (P = 0.26)															

Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias



2 PICO 2 - Ingen medicin + non-farmakologiske interventioner vs waitlist/TAU (uden medicin) (0 RCT)