

ENANNEX I

Identi- fication number of the feed additive	Name of the holder of authori- sation	Name of the additive	Composition, chemical formula, description, analytical method	Species or category of animal	Maxi- mum age	Minimu m content	Maxi mum conten t	Minimu m content	Maxi mum conten t	Other provisions	End of peri- od of auth- orisa- tion
						CFU/kg of complete feed with a moisture content of 12%		CFU/l of water for drinking			
Category: zootechnical additives. Functional group: gut flora stabilisers.											
4b1894	Chr. Hansen A/S	<i>Bacillus subtilis</i> DSM 32324, <i>Bacillus subtilis</i> DSM 32325 and <i>Bacillus</i> <i>amyloliquefaciens</i> DSM 25840	Additive composition Preparation of <i>Bacillus subtilis</i> DSM 32324, <i>Bacillus subtilis</i> DSM 32325 and <i>Bacillus</i> <i>amyloliquefaciens</i> DSM 25840 containing a minimum of 3.2×10^9 CFU/g additive (1.6×10^9 CFU <i>B. subtilis</i> DSM 32324/g; 1.0×10^9 CFU <i>B. subtilis</i> DSM 32325/g and 0.6×10^9 CFU <i>B.</i> <i>amyloliquefaciens</i> DSM 25840/g) (ratio 1.6:1.0:0.6) Solid forms Characterisation of the active substance Viable spores of <i>Bacillus subtilis</i> DSM 32324, <i>Bacillus subtilis</i> DSM 32325 and <i>Bacillus</i>	All poultry species for laying or for breeding	-	1.6×10^9	-	5.4×10^8	-	1. The additive may be used via water for drinking. 2. In the directions for use of the additive and premixtures, the storage conditions, the stability to heat treatment and the stability in water shall be indicated. 3. The additive may be used simultaneously with the following coccidiostats, in accordance with their respective conditions of authorisation as feed additives: diclazuril, decoquinate, halofuginone, monensin, salinomycin, narasin, a combination of nicarbazin and narasin, and lasalocid.	[10 years from the date of entry into force of this Regu- la- tion. To be comp- leted

			<p><i>amyloliquefaciens</i> DSM 25840.</p> <p>Analytical method ⁽¹⁾</p> <p>Identification: DNA sequencing methods or Pulsed Field Gel Electrophoresis (PFGE) (CEN/TS 17697)</p> <p>Enumeration in the feed additive, premixtures, compound feed and water: Spread plate method on tryptone soya agar (EN 15784)</p>							<p>4. For users of the additive and premixtures, feed business operators shall establish operational procedures and organisational measures to address potential risks resulting from their use. Where those risks cannot be eliminated by such procedures and measures, the additive and premixtures shall be used with personal breathing and skin protective equipment.</p>	<p><i>by the Service responsible for the publication]</i></p>
--	--	--	---	--	--	--	--	--	--	--	---

⁽¹⁾ Details of the analytical methods are available at the following address of the Reference Laboratory: https://joint-research-centre.ec.europa.eu/eurl-fa-eurl-feed-additives/eurl-fa-authorisation/eurl-fa-evaluation-reports_en

ANNEX II

Identification number of the feed additive	Name of the holder of authorisation	Name of the additive	Composition, chemical formula, description, analytical method	Species or category of animal	Maximum age	Minimum content	Maximum content	Minimum content	Maximum content	Other provisions	End of period of authorisation
						CFU/kg of complete feed with a moisture content of 12%		CFU/l of water for drinking			
Category: zootechnical additives. Functional group: gut flora stabilisers.											
4b1894	Chr. Hansen A/S	<i>Bacillus subtilis</i> DSM 32324, <i>Bacillus subtilis</i> DSM 32325 and <i>Bacillus amyloliquefaciens</i> DSM 25840	Additive composition Preparation of <i>Bacillus subtilis</i> DSM 32324, <i>Bacillus subtilis</i> DSM 32325 and <i>Bacillus amyloliquefaciens</i> DSM 25840 containing a minimum of 3.2×10^9 CFU/g additive (1.6×10^9 CFU <i>B. subtilis</i> DSM 32324/g; 1.0×10^9 CFU <i>B. subtilis</i> DSM 32325/g and 0.6×10^9 CFU <i>B. amyloliquefaciens</i> DSM 25840/g) (ratio 1.6:1.0:0.6) Solid forms Characterisation of the active substance Viable spores of cells of <i>Bacillus subtilis</i> DSM 32324, <i>Bacillus subtilis</i> DSM 32325 and <i>Bacillus amyloliquefaciens</i> DSM 25840.	All poultry species for fattening or reared for laying or reared for breeding	-	1.6×10^9	-	5.4×10^8	-	1. The additive may be used via water for drinking. 2. In the directions for use of the additive and premixtures, the storage conditions, the stability to heat treatment and the stability in water shall be indicated. 3. The additive may be used simultaneously with the following coccidiostats, in accordance with their respective conditions of authorisation as feed additives: diclazuril, decoquinate, halofuginone, monensin, salinomycin, narasin, a combination of nicarbazin and narasin, and lasalocid.	16.12.2030

			<p><i>Analytical method ⁽²⁾</i></p> <p>Identification: DNA sequencing methods or Pulsed Field Gel Electrophoresis (PFGE) (CEN/TS 17697)</p> <p>Enumeration in the feed additive, premixtures, compound feed and water: Spread plate method on tryptone soya agar (EN 15784).</p>							<p>4. For users of the additive and premixtures, feed business operators shall establish operational procedures and organisational measures to address potential risks resulting from its use. Where those risks cannot be eliminated or reduced to a minimum by such procedures and measures, the additive and premixtures shall be used with personal protective equipment, including breathing protection, glasses and gloves.</p>	
--	--	--	---	--	--	--	--	--	--	---	--

⁽²⁾ Details of the analytical methods are available at the following address of the Reference Laboratory: https://joint-research-centre.ec.europa.eu/eurl-fa-eurl-feed-additives/eurl-fa-authorisation/eurl-fa-evaluation-reports_en